arXiv:math/0701538v2 [math.AG] 19 Mar 2009

RICCI-FLAT DEFORMATIONS OF HOLOMORPHIC VECTOR BUNDLES

MARCO KÜHNEL

Abstract. In this paper we give a criterion for a deformation of a holomorphic vector bundle to be Ricci-flat. As an application we show that on a Kähler manifold, every deformation of a holomorphic vector bundle can be made Ricci-flat whereas on some Hopf manifolds, the non-existence of a Ricci-flat deformation of a vector bundle is connected to non-trivial vector bundles on $\mathbb{C}^n \setminus \{0\}$. In the case of surfaces with $b_1(X) \neq 0$, 'Ricci-rigid' rank 2 vector bundles are proved to be strongly obstructed with respect to possible filtrations. We use this to conclude that on diagonal Hopf surfaces apart from the line bundles only non-good rank 2 bundles can be Ricci-rigid. On Inoue surfaces all Ricci-rigid rank 2 bundles are non-filtrable except one.

1. Introduction

Let X be a compact complex manifold, U a complex manifold without holomorphic vector bundles (e.g. a contractible Stein manifold), and $\mathcal{E} \longrightarrow X \times U$ a holomorphic vector bundle. (So the deformations we deal here with are neither global nor small but something in between.) Further let g be a hermitian metric on $\mathcal{E}$. The data of the central fibre will be denoted by $\mathcal{E}_0$ and g^0. This setup is called a deformation of the hermitian vector bundle $(\mathcal{E}_0, g^0)$. As usual we denote $\Theta^0 := \partial\bar{\partial} \log \det g^0$ and $\Theta^t := \partial_X \overline{\partial}_X \log \det g|_{\mathcal{E}_t}$ the curvatures.

Definition 1.1. *The deformation $(\mathcal{E}, g)$ is called Ricci-flat, if $\Theta^t = \Theta^0$ for all $t \in U$.*

We tackle the question of existence of Ricci-flat deformations. In Theorem 4.1 we show that the obstruction space for this problem is

$$H^1(X, \mathcal{O}_X)/i_* H^1(X, \mathbb{R}),$$

if $i : \mathbb{R} \longrightarrow \mathcal{O}_X$ is the natural inclusion. This group is trivial for Kähler manifolds, so on Kähler manifolds we can extend every metric on the central fibre curvature preservingly to the deformation. In the non-Kähler case we

---

Date: March 20, 2009.

1991 *Mathematics Subject Classification.* 14J60.

Key words and phrases. hermitian, vector bundle, non-filtrable, deformation, Hopf manifold, curvature.

can still show that the property of a given vector bundle to admit only Ricci-flat deformations is independent of the vector bundle and hence a property of the manifold, namely the vanishing of the obstruction space.

The answer to the question, whether for a vector bundle any Ricci-flat deformation is trivial, depends on the bundle, however, and will be denoted by Ricci-rigidity. This generalises the notion of rigidity to manifolds with non-vanishing first cohomology. We will see that this property is connected to a minimizing property of $H^1(\mathcal{E}nd(\mathcal{E}))$.

On Hopf manifolds we deepen this connection and so construct Ricci-rigid vector bundles. We relate Ricci-rigidity to the non-triviality of the pullback to the universal cover $\mathbb{C}^n \setminus \{0\}$.

In the case of a surface with $b_1(X) \neq 0$ Ricci-rigidity of rank 2 bundles will be shown to obstruct possible filtrations. We use this to prove that on an Inoue surface all Ricci-rigid rank 2 bundles are non-filtrable except one (up to tensoring with line bundles) and on Hopf surfaces given by a diagonal automorphism the only Ricci-rigid vector bundles of rank > 1 are of rank 2 and not good in the sense of deformation theory.

It should be noted that the algebraic and analytic category differ widely in this case. Whereas on $\mathbb{C}^2$ and $\mathbb{C}^2 \setminus \{0\}$ there are only trivial algebraic vector bundles [Se58, Ho64], on $\mathbb{C}^2\setminus\{0\}$ some non-trivial holomorphic vector bundles have been constructed before. The paper by Bănică and LePotier [BP87] classified filtrable holomorphic vector bundles on non-algebraic surfaces. In particular, for all integers $r \geq 2, c_2 \geq 0$ there exists a holomorphic rank r bundle $\mathcal{E}$ with $c_2(\mathcal{E}) = c_2$ on a Hopf surface. Calculations below and in [Ma92] show that the pullback to $\mathbb{C}^2 \setminus \{0\}$ of such a bundle is not trivial provided $c_2 > 0$. The constructed bundles are filtrable. Later Ballico [Ba02] constructed a non-filtrable rank 2 bundle on $\mathbb{C}^2 \setminus \{0\}$. In [S66] a non-trivial line bundle on $\mathbb{C}^2 \setminus \{0\}$ is constructed.

2. The local data

By the compactness of X we can employ the trivializations of $\mathcal{E}$ in order to obtain a finite cover of open sets $U_i \subset X$ and isomorphisms

$$\psi_i : \mathcal{E}|_{U_i \times U} \longrightarrow pr_1^*\mathcal{E}_0|_{U_i \times U},$$

where $pr_1 : X \times U \longrightarrow X$ is the projection. This yields the data

$$\theta_{ij} := \psi_i \circ \psi_j^{-1}, g_i := (\psi_i^{-1})^* g,$$

satisfying $g_j = \theta_{ij}^* g_i$, obviously. Note that $\theta_{ij}|_{U_{ij}\times\{0\}} = Id$. Similarly, we have the extended metric of the central fibre

$$<,>:= pr_1^* g^0$$

as a comparison metric. We obtain by Lax-Milgram $G_i \in End_{C^\infty}(pr_1^*\mathcal{E}_0|U_i \times U)$ with the property

$$g_i(e_1, e_2) =< e_1, G_i e_2 >$$

for any C^∞ sections e_1, e_2 of $pr_1^*\mathcal{E}_0$ over U_i. Note that again $G_i|_{U_i \times \{0\}} = Id$. On $U_{ij} \times U$ we have the formula

$$G_i = \theta_{ji}^* G_j \theta_{ji}. \tag{1}$$

Here θ_{ji}^* denotes the adjoint of θ_{ji} with respect to $<,>$. It is easy to see that the deformation is Ricci-flat if and only if

$$\partial_X \overline{\partial}_X \log\det G_i = 0 \text{ for all } i. \tag{2}$$

The inclusions $\mathbb{Z} \subset \mathbb{R} \subset \mathcal{O}_X$ give a commuting triangle

$$\begin{tikzcd}
H^1(X,\mathbb{Z}) \arrow[r] \arrow[dr] & H^1(X,\mathbb{R}) \arrow[d, "i_*"] \\
 & H^1(X,\mathcal{O}_X)
\end{tikzcd}$$

of injective maps. We view all unnamed maps as natural inclusions.

For any deformation the map

$$\eta : U \longrightarrow H^1(X,\mathcal{O}_X)/H^1(X,\mathbb{Z}) = Pic^0(X)$$

given by

$$t \mapsto [\frac{1}{2\pi i}\log\det\theta_{ij}(.,t)] = \det\mathcal{E}_t \otimes \det\mathcal{E}_0^\vee$$

is a well-defined holomorphic map with $\eta(0) = 0$. If we assume the deformation to be small, we may assume that η maps to $H^1(X,\mathcal{O}_X)$.

3. The infinitesimal data

We give the infinitesimal data for the one-dimensional case $U = (\mathbb{C}, 0)$. The generalisation to $U = (\mathbb{C}^n, 0)$ is straightforward.

We take the Taylor series to the first order

$$\theta_{ij} = Id + th_{ij} + h.o.t., G_i = Id + t\gamma_i + \bar{t}\gamma_i^* + h.o.t.$$

with $h_{ij} \in End(\mathcal{E}_0|U_{ij}), \gamma_i \in End_{C^\infty}(\mathcal{E}_0|U_i)$. The h_{ij} satisfy the cocycle condition and hence yield an element $h \in H^1(\mathcal{E}nd(\mathcal{E}_0))$. Indeed, if the deformation is trivial, h_{ij} is a coboundary. By comparison of

$$g_i(e_1, e_2) = < e_1, e_2 > + t < e_1, \gamma_i^* e_2 > + \bar{t} < e_1, \gamma_i e_2 > + h.o.t.$$

and

$$\begin{aligned} \theta_{ji}^* g_j(e_1, e_2) \quad = \quad & < e_1, e_2 > + t(< h_{ji}e_1, e_2 > + < e_1, \gamma_j^* e_2 >) + \\ & + \bar{t} < e_1, (\gamma_j + h_{ji})e_2 > + h.o.t. \end{aligned}$$

we obtain

$$h_{ij} = \gamma_j - \gamma_i \tag{3}$$

on U_{ij}. This tells us that the γ_i trivialise the deformation in the C^∞ sense to the first order.

Now let us consider curvature

$$\begin{aligned}\Theta_i &:= \partial_X\overline{\partial}_X \log\det g_i\\ &= \Theta^0 + \partial\bar\partial \log\det G_i\\ &= \Theta^0 + \partial\bar\partial \log(1 + t\mathrm{tr}\gamma_i + h.o.t)\\ &= \Theta^0 + t\partial\bar\partial \mathrm{tr}\gamma_i + h.o.t.\end{aligned}$$

So, any Ricci-flat deformation satisfies

$$\partial\bar\partial \mathrm{tr}\gamma_i = 0. \tag{4}$$

4. Existence

We start with an arbitrary metric $\tilde g$ on a deformation $\mathcal{E}$. This yields the data $\tilde\psi_i$, $\tilde g^0$ and $\tilde G_i$. By the Cartan decomposition we can find a unique positive hermitian $A \in End_{C^\infty}(\mathcal{E}_0)$ with respect to $\tilde g^0$ such that

$$g^0(e_1, e_2) = \tilde g^0(Ae_1, Ae_2)$$

for any local sections e_1, e_2 of $\mathcal{E}_0$. Setting $\psi_i := A^{-1} \circ \tilde\psi_i, G_i := A^{-1}\tilde G_i A$ (now with respect to $<,>:= pr_1^* g^0$) we obtain a deformation of g^0.

The question of existence of a Ricci-flat deformation is more subtle.

4.1. **Obstructions.**

Theorem 4.1. *A deformation $\mathcal{E} \longrightarrow X \times U$ of a hermitian holomorphic vector bundle $(\mathcal{E}_0 \longrightarrow X, g^0)$ allows for a curvature preserving metric g extending g^0 if and only if*

$$\eta(t) \in i_* H^1(X, \mathbb{R})/H^1(X, \mathbb{Z})$$

for all $t \in U$ and the inclusion $i : \mathbb{R} \hookrightarrow \mathcal{O}_X$; moreover, this condition being satisfied, in every conformal class of metrics g deforming g^0 there is a Ricci-flat representative.

Proof. Necessity: We shrink U_i appropriately such that they are simply connected. By (2) we find for every $t \in U$ holomorphic $h_i(t) \in \mathcal{O}_X(U_i)$ such that

$$\log\det G_i(t) = \mathrm{Re}(h_i(t)).$$

So we have by (1) on U_{ij}

$$2\mathrm{Re}(\log\det\theta_{ij}(t)) = \mathrm{Re}(h_i(t) - h_j(t)),$$

hence there are $c_{ij}(t) \in \mathbb{R}$ such that

$$\frac{1}{2\pi i}(2\log\det\theta_{ij}(t) - h_i(t) + h_j(t)) = c_{ij}(t),$$

but this means exactly

$$\eta(t) \in i_* H^1(X, \mathbb{R})/H^1(X, \mathbb{Z}).$$

Sufficiency: Since $\theta_{ij}|U_{ij} \times \{0\} = Id$, by shrinking U we may assume that $\theta_{ij} = \exp(2\pi i k_{ij})$ for some $k_{ij} \in End(pr_1^*\mathcal{E}_0|U_{ij} \times U)$. Note that (k_{ij}) is no cocycle, unless they commute. Nevertheless, $\det\theta_{ij} = \exp(2\pi i \mathrm{tr} k_{ij})$, so $(\mathrm{tr}k_{ij}) \in \mathcal{O}(U_{ij} \times U)$ is a cocycle, defined uniquely by θ_{ij} up to an integer. In particular, $(\mathrm{tr}k_{ij}|U_{ij} \times \{t\}) \in \mathcal{O}(U_{ij})$ is a cocycle for all $t \in U$. Since we assumed that $\eta(t) = [\mathrm{tr}k_{ij}(t)] \in i_* H^1(X, \mathbb{R})/H^1(X, \mathbb{Z})$, we can choose $K_i(t) \in \mathcal{O}(U_i)$ and $\phi_{ij}(t) \in \mathbb{R}$ such that

$$\mathrm{tr}k_{ij} = K_j - K_i + \phi_{ij}$$

and $K_i(0) = 0, \phi_{ij}(0) = 0$. The dependence on t of K_i and ϕ_{ij} is not holomorphic anymore, but it can be chosen to be C^∞. For $H_i := \exp(2\pi i K_i)$ we obtain

$$\det \theta_{ij} \exp(-2\pi i \phi_{ij}) = \frac{H_j}{H_i}$$

and hence the deformed metric has to satisfy

$$\det G_j = \frac{|H_j|^2}{|H_i|^2} \det G_i. \tag{5}$$

Now we can take any metric $\tilde{g}$ deforming g^0 (with data $\tilde{G}_i$) and take a conformal change:

$$G_i := \left(\frac{|H_i|^2}{\det \tilde{G}_i} \right)^{\frac{1}{n}} \tilde{G}_i.$$

We obtain $\det G_i = |H_i|^2$ and hence $\partial\bar{\partial} \log \det G_i = 0$, so we have a Ricci-flat deformation.

Now covering the original U by neighbourhoods like above and using the uniqueness of $\mathrm{tr}k_{ij}$ up to an integer, we extend the criterion to all of U. ☐

In the infinitesimal information we loose sufficiency of the condition:

Proposition 4.2. *A Ricci-flat deformation germ of hermitian vector bundles satisfies*

$$\frac{1}{2\pi i} trh \in V,$$

where V is the maximal complex subspace of $i_* H^1(X, \mathbb{R})$.

Proof. We shrink U_i appropriately such that they are simply connected. By (4) we find holomorphic $f_i, g_i \in \mathcal{O}(U_i)$ such that $\mathrm{tr}\gamma_i = f_i + \overline{g}_i$. So (3) tells us that

$$\mathrm{tr}h_{ij} + f_i - f_j = \overline{g}_j - \overline{g}_i.$$

Since the right hand side is antiholomorphic and the left hand side is holomorphic, we find constants $c_{ij} \in \mathbb{C}$ such that

$$c_{ij} = \mathrm{tr}h_{ij} + f_i - f_j = \overline{g}_j - \overline{g}_i.$$

So we obtain in $i_* H^1(X, \mathbb{C})$

$$\frac{1}{2\pi i}\mathrm{tr}h = \frac{1}{2\pi i}\mathrm{tr}c = \frac{1}{2\pi i}(\overline{c} + 2i\mathrm{Im}c) = \frac{\mathrm{Im}c}{\pi} \in i_* H^1(X, \mathbb{R}).$$

If we look at a base transformation $\tau : (T,0) \longrightarrow (T,0)$ and denote the objects corresponding to the deformation $\tau^*\mathcal{E}$ of $\mathcal{E}_0$ by a superscript τ, it is straightforward that

$$h_{ij}^{\tau} = \tau'(0)h_{ij}.$$

This proves the statement. □

5. Stable Curvature and Ricci-Rigidity

We see that Ricci-flatness of a deformation does not depend on the initial metric on the central bundle. There are two natural properties connected to Ricci-flat deformations:

Definition 5.1. *The vector bundle $\mathcal{E}_0$ is said to have* stable curvature*, if all deformations of $\mathcal{E}_0$ over a base U without holomorphic vector bundles are Ricci-flat. $\mathcal{E}_0$ is* Ricci-rigid*, if every small Ricci-flat deformation of $\mathcal{E}_0$ is trivial.*

Our first goal is to realise that stable curvature is not a property of a vector bundle, but of the underlying manifold:

Proposition 5.2. *Let X be a compact complex manifold. Then the properties*

- (i) *There exists a vector bundle $\mathcal{E}_0$ with stable curvature,*
- (ii) *All vector bundles have stable curvature,*
- (iii) *$i_* : H^1(X,\mathbb{R}) \longrightarrow H^1(X,\mathcal{O}_X)$ is an $\mathbb{R}$-isomorphism*

are equivalent.

Proof. Since $(ii) \Rightarrow (i)$ is obvious and $(iii) \Rightarrow (ii)$ is Theorem 4.1, it remains to show $(i) \Rightarrow (iii)$. So we assume that i_* is not surjective. In particular, $\dim Pic^0(X) > 0$, hence $\mathcal{O}_X$ can be deformed non-trivially. So, for any deformation $\mathcal{L}_t$ of $\mathcal{L}_0 = \mathcal{O}_X$ with $\mathcal{L}_t \notin i_*H^1(X,\mathbb{R})/H^1(X,\mathbb{Z})$ for all real t and arbitrary vector bundle $\mathcal{E}_0$, the deformation $\mathcal{E}_0 \otimes \mathcal{L}_t$ satisfies $\eta(t) = \mathcal{L}_t^n \notin i_*H^1(X,\mathbb{R})/H^1(X,\mathbb{Z})$ for all real t. Hence we have shown that no vector bundle has stable curvature. □

Due to this result we can define

Definition 5.3. *A compact complex manifold X is called a* stable curvature manifold*, if one (and hence all) of the properties in Proposition 5.2 are satisfied.*

Examples of stable curvature manifolds are all compact Kähler manifolds. Counterexamples are all compact manifolds with odd $b_1(X)$, e.g. Hopf surfaces.

Also Ricci-rigidity has geometric implications.

Proposition 5.4. *If X allows for a Ricci-rigid vector bundle, then all germs of holomorphic maps $f : (\mathbb{C},0) \longrightarrow Pic^0(X)$ with image in*

$$i_*H^1(X,\mathbb{R})/H^1(X,\mathbb{Z})$$

are constant; in particular, Ricci-flat deformations are exactly the ones with preserved determinant; and $V = \{0\}$.

Proof. Let f be such a map with $f(0) = 0$ and $\mathcal{E}_0$ a Ricci-rigid vector bundle. Then, of course $\mathcal{E}_0 \otimes f(t)$ is a Ricci-flat deformation, hence $\mathcal{E}_0 \otimes f(t) \cong \mathcal{E}_0$. Taking determinants shows $f(t)^n = \mathcal{O}_X$, but this means that $f(t)$ is constant, so $f \equiv 0$. $\square$

Note, that for any vector bundle the sequence

$$0 \longrightarrow \mathcal{O}_X \xrightarrow{\cdot Id} \mathcal{E}nd(\mathcal{E}_0) \xrightarrow{\psi} ad(\mathcal{E}_0) \longrightarrow 0$$

splits via the trace map:

$$\mathcal{E}nd(\mathcal{E}_0) \cong \mathcal{O}_X \oplus ad(\mathcal{E}_0), \phi \mapsto \left(\frac{\mathrm{tr}\phi}{\mathrm{rk}\,\mathcal{E}_0}, \psi(\phi)\right)$$

is an isomorphism.

In particular, $tr : H^1(\mathcal{E}nd(\mathcal{E}_0)) \longrightarrow H^1(\mathcal{O}_X)$ is surjective. If $\mathcal{E}_0$ is good, i.e. $H^2(ad(\mathcal{E}_0)) = 0$, then we can integrate any holomorphic $h : U \longrightarrow V$ on a small open set $U \subset \mathbb{C}$ to a Ricci-flat deformation of $\mathcal{E}_0$.

The classical correspondence that a vector bundle $\mathcal{E}_0$ over X with $H^2(ad(\mathcal{E}_0)) = 0$ is rigid if and only if $H^1(X, \mathcal{E}nd(\mathcal{E}_0)) = 0$ has a nice analogue here. Our notion of Ricci-rigidity measures in the infinitesimal setup that at $\mathcal{E}_0$ the minimal value is attained.

Proposition 5.5. *If a good vector bundle* $\mathcal{E}_0$ *on* X *is Ricci-rigid, then* $H^1(ad(\mathcal{E}_0)) = 0$.

Proof. Let $\mathcal{E}_0$ be Ricci-rigid and $\mathcal{E}_t$ be any small deformation of $\mathcal{E}_0$ and define $\tilde{\mathcal{L}}_t := \det \mathcal{E}_t^\vee \otimes \det \mathcal{E}_0$. Note that $\tilde{\mathcal{L}}_t \in Pic^0(X)$. If we shrink the base of the deformation appropriately we may assume the existence of a holomorphic family $\mathcal{L}_t$ of line bundles with $\mathcal{L}_t^n = \tilde{\mathcal{L}}_t$. Now we have

$$\det(\mathcal{E}_t \otimes \mathcal{L}_t) = \det(\mathcal{E}_0),$$

in particular, $\mathcal{E}_t \otimes \mathcal{L}_t$ is a Ricci-flat deformation. Hence we obtain

$$\mathcal{E}_t \cong \mathcal{E}_0 \otimes \mathcal{L}_t^\vee.$$

By imposing $H^2(\mathcal{E}nd(\mathcal{E}_0)) = H^2(\mathcal{O}_X)$ we ensure that every $\zeta \in H^1(\mathcal{E}nd(\mathcal{E}_0))$ can be integrated to a small deformation (see [Br95, Cor. 5.7]). So

$$\zeta = \frac{d}{dt}|_{t=0}\mathcal{L}_t{}^\vee \in H^1(\mathcal{O}_X)$$

for some deformation $\mathcal{L}_t \in Pic^0(X)$ of $\mathcal{L}_0 = \mathcal{O}_X$. $\square$

Proposition 5.6. *Let* X *be a compact manifold allowing for a Ricci-rigid vector bundle and* $\mathcal{E}_0$ *a holomorphic vector bundle on* X. *If* $H^1(ad(\mathcal{E}_0)) = 0$, *then* $\mathcal{E}_0$ *is Ricci-rigid.*

Proof. If $H^1(\mathcal{E}nd(\mathcal{E}_0)) = H^1(\mathcal{O}_X)$, and $\mathcal{E}_t$ is a Ricci-flat deformation, then by Grauert's semi-continuity theorem we obtain also $H^1(\mathcal{E}nd(\mathcal{E}_t)) = H^1(\mathcal{O}_X)$ for small t. Shifting the centre of the deformation to t, we obtain a family $h(t) \in H^1(\mathcal{E}nd(\mathcal{E}_t))$. We know now that $\mathrm{tr} : H^1(\mathcal{E}nd(\mathcal{E}_t)) \longrightarrow H^1(\mathcal{O}_X)$ is an isomorphism. By the Ricci-flatness of the deformation we have due to Proposition 5.4 that $\mathrm{tr}h(t) = 0$. Hence $h(t) = 0$ for all t and so the deformation is trivial. □

6. Examples

Since all Kähler manifolds are stable curvature manifolds, Ricci-rigidity coincides with rigidity in these cases. Therefore we concentrate on non-Kähler examples.

6.1. **Hopf manifolds.**

Example 6.1. Let X be the Hopf manifold defined by the automorphism $\phi(z) = 2z$ on $\mathbb{C}^n \setminus \{0\}$. Then there is a natural smooth elliptic fibration $\pi : X \longrightarrow \mathbb{P}^{n-1}$. For any bundle we have $R^1\pi_*\pi^*\mathcal{E}_0 = \mathcal{E}_0^\vee$. Let $\mathcal{E}_0$ be a a simple bundle on $\mathbb{P}^{n-1}$ with $H^1(\mathcal{E}nd(\mathcal{E}_0)) = H^2(\mathcal{E}nd(\mathcal{E}_0)) = 0$. Then the Leray spectral sequence implies that $H^1(\mathcal{E}nd(\pi^*\mathcal{E}_0)) = \mathbb{C}$ and hence $\pi^*\mathcal{E}_0$ is Ricci-rigid. For instance, $T_{\mathbb{P}^{n-1}}$ satisfies these conditions for $n \geq 2$. (For $n = 2$ exactly the line bundles satisfy the requirements.) △

Moreover, it is known that $p^*T_{\mathbb{P}^n}$ is not trivial for $n \geq 2$, if $p : \mathbb{C}^{n+1} \setminus \{0\} \longrightarrow \mathbb{P}^n$ denotes the natural projection. The following results will recover this and give a connection between Ricci-rigid bundles on some Hopf manifolds and non-trivial vector bundles on $\mathbb{C}^n \setminus \{0\}$.

Proposition 6.2. *Let X be the Hopf manifold given by the quotient of $\mathbb{C}^n \setminus \{0\}$ by the automorphism group generated by $\varphi(z_1, \ldots, , z_n) = (\alpha_1 z_1, \ldots, \alpha_n z_n)$, $|\alpha_i| > 1$ and $u : \mathbb{C}^n \setminus \{0\} \longrightarrow X$ the projection. If $\mathcal{E}_0$ is a Ricci-rigid vector bundle on X with $\mathrm{rk}\,\mathcal{E}_0 > 1$, then $u^*\mathcal{E}_0$ is not trivial.*

Proof. There is a multiplicative degree $\deg_\varphi : \mathbb{C}[z_1, \ldots, z_n] \longrightarrow \mathbb{C}$ via

$$\deg_\varphi(z_i) := \alpha_i.$$

If $u^*\mathcal{E}_0$ is trivial, usual techniques allow us to identify $\mathcal{E}_0$ of rank r with an equivalence class of holomorphic maps $L : \mathbb{C}^n \setminus \{0\} \longrightarrow Gl(r, \mathbb{C})$ where

$$L \cong \tilde{L} :\Longleftrightarrow \quad \exists T \in \mathcal{O}(\mathbb{C}^n \setminus \{0\}, Gl(r, \mathbb{C})) \text{ such that } \tilde{L} = T \circ \varphi \cdot L \cdot T^{-1}.$$

Choosing T carefully we can achieve a normal form of L consisting of blocks $L_\nu, \nu \in \mathbb{C}^*$ in upper triangle form (cf. [Ma92] for a very similar normal form) with the property

$$(L_\nu)_{kk} = \prod_{j=1}^{n} \alpha_j^{i_{jk}} \nu$$

for $i_{jk} \geq 0, i_{j1} = 0$ and

$$(L_\nu)_{kl} \in \mathbb{C}[z_1, \dots, z_n]$$

homogeneous with

$$\deg_\varphi (L_\nu)_{kl} = \prod_{j=1}^{n} \alpha_j^{i_{jl} - i_{jk}}.$$

Now $L(t)_{kl} := L_{kl}$ for $(k, l) \notin \{(1,1), (2,2)\}$ and

$$L(t)_{11} := \exp(t) L_{11}, L(t)_{22} := \exp(-t) L_{22}$$

defines a non-trivial Ricci-flat small deformation of $\mathcal{E}_0$. □

Combining (6.2) and (6.1) we obtain immediately

Corollary 6.3. *If* $n > 1, p : \mathbb{C}^{n+1} \setminus \{0\} \longrightarrow \mathbb{P}^n$ *is the natural projection and* $\mathcal{E}_0$ *a simple vector bundle on* $\mathbb{P}^n$ *satisfying* $\operatorname{rk} \mathcal{E}_0 > 1$ *and* $H^1(\mathcal{E}nd(\mathcal{E}_0)) = H^2(\mathcal{E}nd(\mathcal{E}_0)) = 0$*, then* $p^* \mathcal{E}_0$ *is not trivial.*

6.2. **Surfaces.**

Theorem 6.4. *Let* X *be a compact complex surface with* $b_1(X) \neq 0$*. If a rank 2 vector bundle* $\mathcal{E}_0$ *on* X *is Ricci-rigid, then every filtration*

$$0 \longrightarrow \mathcal{F} \longrightarrow \mathcal{E}_0 \longrightarrow \mathcal{G} \otimes I_Z \longrightarrow 0$$

with Z *a (non-reduced) point set and line bundles* $\mathcal{F}, \mathcal{G}$ *satisfies* $H^1(X, \mathcal{F} \otimes \mathcal{G}^\vee) \neq 0$.

Proof. We may assume that X is not Kähler; otherwise there are no Ricci-rigid vector bundles. Assume we have an extension

$$0 \longrightarrow \mathcal{F} \longrightarrow \mathcal{E}_0 \longrightarrow \mathcal{G} \otimes I_Z \longrightarrow 0$$

with line bundles $\mathcal{F}, \mathcal{G}$ and a finite set of points Z such that $H^1(X, \mathcal{F} \otimes \mathcal{G}^\vee) = 0$.

This extension yields an element $\zeta \in \operatorname{Ext}^1(I_z, \mathcal{F} \otimes \mathcal{G}^\vee)$. A part of the long exact *Ext*-sequence of the ideal sheaf sequence looks like

$$0 = H^1(X, \mathcal{F} \otimes \mathcal{G}^\vee) \longrightarrow \operatorname{Ext}^1(I_Z, \mathcal{F} \otimes \mathcal{G}^\vee) \longrightarrow \operatorname{Ext}^2(\mathcal{O}_Z, \mathcal{F} \otimes \mathcal{G}^\vee) \longrightarrow H^2(X, \mathcal{F} \otimes \mathcal{G}^\vee).$$

So ζ may be identified with an element $\xi \in \operatorname{Ext}^2(\mathcal{O}_Z, \mathcal{G}^\vee \otimes \mathcal{F})$ mapping to $0 \in H^2(X, \mathcal{G}^\vee \otimes \mathcal{F})$ and generating $\mathcal{E}xt^2(\mathcal{O}_Z, \mathcal{F} \otimes \mathcal{G}^\vee) \cong \mathcal{O}_Z$. Now let $\mathcal{L}_t$ be a non-constant holomorphic family of line bundles with $\mathcal{L}_0 = \mathcal{O}_X$; due to $b_1(X) \neq 0$ such a family exists. We interpret

$$\xi \in \operatorname{Ext}^2(\mathcal{O}_Z, \mathcal{G}^\vee \otimes \mathcal{F}) \cong \operatorname{Ext}^2(\mathcal{O}_Z \otimes \mathcal{L}_t^{-2}, \mathcal{G}^\vee \otimes \mathcal{F}) \cong \operatorname{Ext}^2(\mathcal{O}_Z, \mathcal{G}^\vee \otimes \mathcal{F} \otimes \mathcal{L}_t^2),$$

again generating $\mathcal{E}xt^2(\mathcal{O}_Z, \mathcal{F} \otimes \mathcal{G}^\vee \otimes \mathcal{L}_t^2) \cong \mathcal{O}_Z$. By Grauert's semicontinuity and Lemma 6.5 we may assume

$$H^1(X, \mathcal{F} \otimes \mathcal{G}^\vee \otimes \mathcal{L}_t^2) = 0$$

for all t small enough and

$$H^2(X, \mathcal{F} \otimes \mathcal{G}^\vee \otimes L_t^2) = 0$$

for all $t \neq 0$ small enough. So by the analogue *Ext*-sequence argument we obtain a unique $\zeta_t \in \operatorname{Ext}^1(I_Z, \mathcal{F} \otimes \mathcal{G}^\vee \otimes \mathcal{L}_t^2)$, i.e. we obtain by the Serre construction (see e.g. [Br95]) an extension

$$0 \longrightarrow \mathcal{F} \otimes \mathcal{L}_t \longrightarrow \mathcal{E}_t \longrightarrow \mathcal{G} \otimes \mathcal{L}_t^\vee \otimes I_Z \longrightarrow 0$$

with a vector bundle $\mathcal{E}_t$. The Ricci-flat deformation of $\mathcal{E}_0$ obtained in this way is not trivial, again due to Lemma 6.5: Of course, $H^0(\mathcal{E}_0 \otimes \mathcal{F}^\vee) \neq 0$ and $H^0(\mathcal{E}_t \otimes \mathcal{F}^\vee \otimes \mathcal{L}_t^\vee) \neq 0$, hence $H^0(\mathcal{E}_t \otimes \mathcal{F}^\vee) = 0$ for general t, if the family $\mathcal{L}_t$ was chosen general enough. □

Lemma 6.5. *On a compact, non-Kähler surface X for every line bundle $\mathcal{F}$ the set*

$$\{\mathcal{L} \in Pic^0(X) |\ H^0(\mathcal{F} \otimes \mathcal{L}) = 0\}$$

is open and not empty. The same holds true, if $\mathcal{F}$ is a filtrable rank 2 vector bundle.

Proof. First, let $\mathcal{F}$ be a line bundle. By Grauert's semicontinuity result it is enough to find some $\mathcal{L} \in Pic^0(X)$ with $H^0(\mathcal{F} \otimes \mathcal{L}) = 0$. So we assume that $H^0(\mathcal{F} \otimes \mathcal{L}) \neq 0$ for all $\mathcal{L} \in Pic^0(X)$. Note that due to the non-Kähler property $c_1(\mathcal{G})^2 \leq 0$ for every line bundle $\mathcal{G}$ (cf. [BHPV, IV,Thm 6.2]) and $b_1(X)$ is odd (cf. [BHPV, IV,Thm 3.1]). The latter implies that $Pic^0(X)$ is not compact.

In a first step we construct a fibration $\pi : X \longrightarrow C$. For $s \in H^0(\mathcal{F})$ and $t \in H^0(\mathcal{F} \otimes \mathcal{L})$ for some $\mathcal{L} \in Pic^0(X)$ we see that due to

$$c_1(\mathcal{F}).c_1(\mathcal{F} \otimes \mathcal{L}) = c_1(\mathcal{F})^2 \leq 0$$

the schemes $V(s), V(t)$ do not intersect or have at least one common component. So

$$D := \bigcap_{\mathcal{L} \in Pic^0(X)} V(H^0(\mathcal{F} \otimes \mathcal{L}))$$

is empty or an effective divisor. The line bundle

$$\mathcal{G} := \mathcal{F} \otimes \mathcal{O}_X(-D)$$

again satisfies $H^0(\mathcal{G} \otimes \mathcal{L}) \neq 0$ for all $\mathcal{L} \in Pic^0(X)$ and, moreover, we can find $\mathcal{L} \in Pic^0(X)$ and sections $s_0 \in H^0(\mathcal{G})$, $s_1 \in H^0(\mathcal{G} \otimes \mathcal{L})$ and $s_{-1} \in H^0(\mathcal{G} \otimes \mathcal{L}^\vee)$ such that

$$V(s_0) \cap V(s_1) = V(s_0) \cap V(s_{-1}) = \emptyset.$$

Hence,

$$\pi' : X \longrightarrow \mathbb{P}^1, x \mapsto [s_0^2 : s_1 s_{-1}]$$

is a well-defined map. Applying Stein factorization we obtain a fibration

$$\pi : X \longrightarrow B$$

onto a smooth curve B with connected fibres.

Note that all irreducible curves $C \subset X$ are contained in the fibres, since otherwise we have a curve C with $C.F > 0$ for the general fibre F of π; but $(C + kF)^2 = C^2 + 2kC.F > 0$ for k big enough.

Let $C_1, \ldots, C_n$ be the finitely many irreducible components of the reducible fibres of π. Now we have proved that for the compact manifold $M := B \cup \{C_1, \ldots, C_n\}$ and $P := \mathcal{F} \otimes Pic^0(X)$, the connected component of $Pic(X)$ containing $\mathcal{F}$, we have a map

$$\psi : \bigcup_{k \in \mathbb{N}_0} M^k \longrightarrow Pic(X)$$

induced by $\psi(b) := \mathcal{O}_X(F_b)$ for $b \in B$ and $\psi(C_i) := \mathcal{O}_X(C_i)$ satisfying

$$P \subset \mathrm{im}\psi.$$

On the other hand $\psi(M^k)$ is compact and hence of codimension at least 1 in $Pic(X)$. So $\mathrm{im}\psi$ cannot contain any connected component of $Pic(X)$ and we have reached a contradiction.

If $\mathcal{F}$ is a filtrable rank 2 bundle, we take a filtration

$$0 \longrightarrow \mathcal{F}' \longrightarrow \mathcal{F} \longrightarrow \mathcal{F}'' \otimes I_Z \longrightarrow 0$$

and apply the lemma to the line bundles $\mathcal{F}'$ and $\mathcal{F}''$. ☐

6.2.1. *Ricci-rigid rank* 2 *bundles on Inoue surfaces.* We treat Inoue surfaces before Hopf surfaces, since the description of Ricci-rigid vector bundles is somewhat easier due to the lack of curves. The geometry of Inoue surfaces implies immediately that all line bundles are Ricci-rigid. So we tackle rank 2 bundles.

Corollary 6.6. *On an Inoue surface X every Ricci-rigid rank 2 vector bundle $\mathcal{E}_0$ is either non-filtrable or there is a line bundle $\mathcal{F}$ such that $\mathcal{E}_0 \otimes \mathcal{F}$ is isomorphic to the unique vector bundle obtained as a non-split extension*

$$0 \longrightarrow K_X \longrightarrow \mathcal{E}_0 \otimes \mathcal{F} \longrightarrow \mathcal{O}_X \longrightarrow 0.$$

The latter is Ricci-rigid, indeed.

Proof. By tensoring with line bundles we may assume that $\mathcal{E}_0$ allows for a filtration

$$0 \longrightarrow \mathcal{L} \longrightarrow \mathcal{E}_0 \longrightarrow I_Z \longrightarrow 0.$$

By Theorem 6.4 we obtain $H^1(X, \mathcal{L}) \neq 0$. On an Inoue surface the intersection form of line bundles is trivial, hence $\chi(\mathcal{L}) = \chi(\mathcal{O}_X) = 0$; furthermore

$$H^0(\mathcal{F}) \neq 0 \iff \mathcal{F} = \mathcal{O}_X$$

for any line bundle, since there are no curves on X. So the condition $H^1(X, \mathcal{L}) \neq 0$ translates into $\mathcal{L} \in \{K_X, \mathcal{O}_X\}$.

First we treat the case $\mathcal{L} = K_X$ and $Z \neq \emptyset$. The filtration dualises to

$$0 \longrightarrow \mathcal{O}_X \longrightarrow \mathcal{E}_0^\vee \longrightarrow K_X^\vee \otimes I_Z \longrightarrow 0.$$

So $H^0(X, \mathcal{E}_0^\vee) = \mathbb{C}$ and, looking at the dualised sequence again, also $H^0(X, \mathcal{E}_0^\vee \otimes I_Z) = \mathbb{C}$. If we tensorise this sequence by $\mathcal{E}_0$ we obtain

$$0 \longrightarrow \mathcal{E}_0 \longrightarrow \mathcal{E}nd(\mathcal{E}_0) \longrightarrow \mathcal{E}_0^\vee \otimes I_Z \longrightarrow 0,$$

yielding for global sections $H^0(X, \mathcal{E}nd(\mathcal{E}_0)) = \mathbb{C}$, so $\mathcal{E}_0$ is simple. Tensoring the filtration and its dual with K_X tells us $H^0(X, \mathcal{E}_0 \otimes K_X) = 0 = H^0(X, \mathcal{E}_0^\vee \otimes K_X \otimes I_Z)$, so also

$$H^2(X, \mathcal{E}nd(\mathcal{E}_0) = H^0(X, \mathcal{E}nd(\mathcal{E}_0) \otimes K_Z) = 0.$$

Now we employ

$$-4c_2(\mathcal{E}_0) = \chi(\mathcal{E}nd(\mathcal{E}_0)) = 1 - h^1(X, \mathcal{E}nd(\mathcal{E}_0)).$$

Theorem 5.5 implies that $\mathcal{E}_0$ is not Ricci-rigid.

If $Z = \emptyset$ the filtration is

$$0 \longrightarrow K_X \longrightarrow \mathcal{E}_0 \longrightarrow \mathcal{O}_X \longrightarrow 0.$$

If the sequence splits, $\mathcal{E}_0$ is clearly not Ricci-rigid. For the non-splittung extension we have $H^0(\mathcal{E}_0) = 0$. The dual of the filtration implies $H^0(\mathcal{E}_0^\vee) = \mathbb{C}$, so like above we obtain that $\mathcal{E}_0$ is simple and $H^2(X, \mathcal{E}nd(\mathcal{E}_0)) = 0$. Riemann-Roch implies this time $H^1(X, \mathcal{E}nd(\mathcal{E}_0)) = \mathbb{C}$ and so $\mathcal{E}_0$ is Ricci-rigid, indeed.

We are left with the case $L = \mathcal{O}_X$. Now $\mathcal{E}_0 \cong \mathcal{E}_0^\vee$ and with the same techniques as above we see that $H^2(X, \mathcal{E}nd(\mathcal{E}_0)) = 0$ and $h^0(\mathcal{E}nd(\mathcal{E}_0)) \in \{1, 2\}$, if $Z \neq \emptyset$. Riemann-Roch yields again that $\mathcal{E}_0$ is not Ricci-rigid. If $Z = \emptyset$, the non-splitting extension

$$0 \longrightarrow \mathcal{O}_X \longrightarrow \mathcal{E}_0 \longrightarrow \mathcal{O}_X \longrightarrow 0$$

satisfies $H^0(X, \mathcal{E}_0) = \mathbb{C}$. Hence $h^0(X, \mathcal{E}nd(\mathcal{E}_0)) \geq 2$. Moreover,

$$H^1(X, \mathcal{E}_0) = \mathbb{C}, H^2(X, \mathcal{E}_0) = 0.$$

The sequence

$$0 \longrightarrow \mathcal{E}_0 \longrightarrow \mathcal{E}nd(\mathcal{E}_0) \longrightarrow \mathcal{E}_0 \longrightarrow 0$$

now implies $H^2(X, \mathcal{E}nd(\mathcal{E}_0)) = 0, h^0(X, \mathcal{E}nd(\mathcal{E}_0)) = h^1(X, \mathcal{E}nd(\mathcal{E}_0)) = 2$, so $\mathcal{E}_0$ is not Ricci-rigid. ☐

6.2.2. *Ricci-rigid vector bundles on Hopf surfaces.*

Corollary 6.7. *Let X be a Hopf surface given by diagonal φ. All line bundles on X are Ricci-rigid. There are no Ricci-rigid vector bundles of rank ≥ 3. There are no good Ricci-rigid rank 2 vector bundles.*

Proof. By Proposition 5.6 the line bundles are Ricci-rigid.

Next we observe that a vector bundle $\mathcal{G}$ on $\mathbb{C}^2 \setminus \{0\}$ is trivial if and only if it is extendable to $\mathbb{C}^2$ (see [S66]). It is extendable as a vector bundle if and only if it is extendable as a coherent sheaf, but this is ensured by [FG69, Cor. VII.4], if $\operatorname{rk} \mathcal{G} \geq 3$. Hence, by Proposition 6.2, a vector bundle $\mathcal{E}_0$ on a Hopf surface given by diagonal φ is never Ricci-rigid, if $\operatorname{rk} \mathcal{E}_0 > 2$. So it remains to deal with the "critical case" $\operatorname{rk} \mathcal{E}_0 = 2$.

In case $\mathcal{E}_0$ is Ricci-rigid, good and $\operatorname{rk} \mathcal{E}_0 = 2$ we have $H^1(X, \mathcal{E}nd(\mathcal{E}_0)) = \mathbb{C}$ and deduce

$$-4c_2(\mathcal{E}_0) = \chi(\mathcal{E}nd(\mathcal{E}_0)) \geq 0.$$

If $c_2(\mathcal{E}_0) < 0$, then $\mathcal{E}_0$ is not filtrable, so it is simple; but then $c_2(\mathcal{E}_0) = 0$. If $c_2(\mathcal{E}_0) = 0$, then $\mathcal{E}_0$ is simple and topologically trivial, contradicting [Mo04, Prop 3.1]. □

References

[AT03] Aprodu, M., Toma, M., 'Une note sur les fibrés holomorphes non-filtrables. (French. English, French summary) [A note on non-filtrable holomorphic bundles]', *C. R. Math. Acad. Sci. Paris* 336(7), 581–584 (2003)

[Ba02] Ballico, E., 'Holomorphic vector bundles on $\mathbb{C}^2 \setminus \{0\}$', *Isr. J. of Math.* 128, 197–204 (2002)

[BP87] Bănică, C., Le Potier, J., 'Sur l'existence des fibrs vectoriels holomorphes sur les surfaces non-algbriques', *J. Reine Angew. Math.* 378, 1–31 (1987)

[BHPV] Barth, W., Hulek, K., Peters, C., van de Ven, A., 'Compact Complex Surfaces', Springer, 2004

[Br95] Brînzănescu, V., 'Holomorphic Vector Bundles over Compact Complex Surfaces', Springer, 1996

[BrMo05] Brînzănescu, V., Moraru, R., 'Stable Bundles on Non-Kähler Elliptic Surfaces', *Commun. Math. Phys.* 254, 565–580 (2005)

[FG69] Frisch, J.; Guenot, J., 'Prolongement de faisceaux analytiques cohérents', *Invent. Math.* 7, 321–343 (1969)

[Ha] Hartshorne, R., 'Algebraic Geometry', Springer, 1977

[HS] Hilton, P.J., Stammbach, U., 'A Course in Homological Algebra', Springer, 1971

[Ho64] Horrocks, G., 'Vector bundles on the punctured spectrum of a local ring', *Proc. London Math. Soc. (3)* 14, 689–713 (1964)

[Mo04] Moraru, R., 'Stable bundles on Hopf manifolds', math.AG/0408439 (2004)

[Ma92] Mall, D., 'On holomorphic vector bundles on Hopf manifolds with trivial pullback on the universal covering', *Math. Ann.* 294, 718–740 (1992)

[S66] Serre, J.-P., 'Prolongement de faisceaux analytiques cohérents', *Ann. Inst. Fourier* 16, No.1, 363–374 (1966)

[Se58] Seshadri, C.S., 'Triviality of vector bundles over the affine space K^2', *Proc. Natl. Acad. Sci. USA* 44, 456–458 (1958)

[Ve06] Verbitsky, M., 'Holomorphic bundles on diagonal Hopf manifolds', *Izvestiya* 70(5), 867–882 (2006)

Marco Kühnel, Otto-von-Guericke-Universität Magdeburg, FMA / IAN, Postfach 4120, 39016 Magdeburg, Germany

E-mail address: `marco.kuehnel@ovgu.de`